

## Obstetrics & Gynecology:

January 2012 - Volume 119 - Issue 1 - p 14–20

doi: 10.1097/AOG.0b013e31823d4167

Original Research

# Effect of Daily Text Messages on Oral Contraceptive Continuation: A Randomized Controlled Trial

**Castaño, Paula M. MD, MPH; Bynum, Jillian Y. MPH; Andrés, Raquel MA, PhD; Lara, Marcos PharmD, MBA; Westhoff, Carolyn MD, MSc**



### Author Information

From the Department of Obstetrics and Gynecology, Columbia University Medical Center, New York, New York.

Funded in part by Affinity Health Plan Making a World of Difference Grant Program. Additional funding was provided by the William and Flora Hewlett Foundation.

The authors thank Planned Parenthood of New York City for providing the clinical site (Planned Parenthood Boro Hall) and ConnectUS and MIR3 for sending the text messages.

Preliminary results were presented at Reproductive Health 2010, September 22–25, 2010, Atlanta, Georgia. Additional results were presented at The North American Forum on Family Planning, October 22–24, 2011, Washington, DC.

The opinions expressed in this article do not necessarily reflect those of Planned Parenthood Federation of America, Inc.

Corresponding author: Paula M. Castaño, MD, MPH, Department of Obstetrics and Gynecology, Columbia University Medical Center, 622 W 168th Street, PH 16, New York, NY 10032; e-mail: pc2137@columbia.edu.

**Financial Disclosure** Dr. Castaño serves on the scientific advisory board for Bayer. Dr. Westhoff is an advisory board member for Teva and Agile and is a consultant for Merck, Bayer, and Med360. The other authors did not report any potential conflicts of interest.



**Abstract**

**OBJECTIVE:** To estimate whether daily educational text messages affect oral contraceptive pill (OCP) continuation at 6 months.

**METHODS:** We randomized young women electing OCPs at an urban family planning health center to either routine care or routine care plus 180 days of daily educational text messages. Investigators masked to treatment allocation randomized participants who were not masked to treatment. The primary outcome measure was self-reported OCP continuation through a telephone call at 6 months (contacts between 5 and 8 months).

**RESULTS:** We enrolled 962 participants (480 intervention and 482 routine care) and obtained continuation data on 683 (346 and 337, respectively). At the follow-up, 64% of participants randomized to the intervention were still OCP users compared with 54% of the routine care group ( $P=.005$ ). Continuation was highest in the intervention group if the interview took place while the intervention was ongoing (75% compared with 54%,  $P=.003$ ); the effect of the intervention on continuation was less after the intervention ended (60% compared with 54%,  $P=.16$ ). Participants receiving the intervention were more likely to continue oral contraception than control participants at 6 months (odds ratio 1.44, 95% confidence interval 1.03–2.00) in analyses adjusted for age, race or ethnicity, age at coitarche, pregnancy history, and OCP experience.

**CONCLUSION:** The use of daily educational text messages improves OCP continuation at 6 months over routine care alone. Ten women would need to receive this simple intervention to improve continuation in one. This effect is strongest in the women whose follow-up took place while the text intervention was ongoing.

**CLINICAL TRIAL REGISTRATION:** ClinicalTrials.gov, [www.ClinicalTrials.gov](http://www.ClinicalTrials.gov), NCT00677703.

**LEVEL OF EVIDENCE:** I

Fifty percent of patients worldwide take medications improperly, and this poor adherence to medications can result in adverse outcomes.<sup>1</sup> Young women are most likely to choose oral contraceptive pills (OCPs) for pregnancy prevention,<sup>2</sup> but improper use and discontinuation are common. Six-month OCP continuation rates in young women range from 12% to 58%.<sup>3–5</sup> Oral contraceptive pill misuse, failure, and discontinuation lead to 1 million unintended pregnancies each year in the United States.<sup>6</sup> Failure to establish a routine of taking the pill is a common reason for OCP discontinuation.<sup>7</sup>

Text messaging has been investigated as an adjunct to routine health care, including the management of chronic medical conditions and medication regimen adherence. Studies have evaluated text messages as behavior modifiers,<sup>8,9</sup> laboratory result services,<sup>10</sup> and

medication reminders.<sup>11,12</sup> Only one trial has evaluated the use of daily text messages on OCP adherence and yielded null results. In that study of 82 women, both the control and the intervention groups used an electronic device that tracked pill-taking behavior for 3 months. Use of the device did not resemble ordinary OCP use; thus, it is difficult to interpret the results. That study did confirm the prevalence of missed OCPs with all participants missing nearly five pills per cycle.<sup>13</sup>

Young women lead the way in cell phone use.<sup>14</sup> Monthly cell phone bills reveal that women under age 18 years send and receive 4,050 text messages monthly followed by young adult women up to age 24 years who send and receive 1,630 text messages monthly.<sup>15</sup> Our study estimates whether a daily educational text message, in addition to routine clinical care, affects 6-month OCP continuation rates in young women.

## **MATERIALS AND METHODS**

We recruited women younger than age 25 years at a Planned Parenthood family planning health center in downtown Brooklyn, New York. Clients were eligible for participation if they were sexually active, owned a cell phone with text messaging functionality, and had no medical contraindications to oral contraceptive use. The Columbia University Medical Center institutional review board and the Research Department at Planned Parenthood Federation of America approved the study. Bilingual study staff screened all participants, who provided written informed consent, completed a baseline questionnaire, and agreed to be contacted by phone after 6 months for an exit questionnaire. At enrollment, we provided participants with \$10 compensation, the average cost of 180 additional text messages.

We randomized participants to receive routine care or routine care plus a daily text message for 6 months using 1:1 fixed allocation randomization with blocks of four. Staff members who had no contact with the enrollment site or study participants used a random-number table to generate the sequence and placed assignments into sequentially numbered, sealed opaque envelopes. We kept the sequence in a password-protected database. Randomization envelopes were opened by recruiters at the enrollment site after all enrollment procedures were completed. Study participants could not be masked.

All participants received routine care following health center protocols, including contraceptive counseling by staff and an educational information handout detailing use, effectiveness, benefits, and risks. The intervention group also received 180 daily text messages that included an introductory message, three reminders of how to change contact information or message time, 47 individual educational messages that were repeated up to four times, 12 two-way messages for quality control, and a final message (available in the Appendix online at <http://links.lww.com/AOG/A272>). The educational messages were adapted from information in the handout, were both positive and negative in content, and included: "The pill improves anemia" and "Women who use the pill have a slightly greater risk of developing blood clots in their legs." The educational messages incorporated six domains of OCP knowledge: risks, benefits, side effects, use, effectiveness, and mechanisms

of action.<sup>16</sup> The two-way messages asked for participant feedback and included: “We want to know if you are receiving this message at the time you requested. Please reply 1 for Yes or 2 for No.” We maintained a total message length under 160 characters. To maintain confidentiality, the messages did not refer to the participant by name or specify that they were reminders to take OCPs. We prepared Spanish research documents and text messages, but all participants chose to complete documents and receive messages in English.

The notification company began to send out text messages using a secure web-based platform on a start date specified by the participant at enrollment. The start date depended on the participant's OCP use status (new start compared with ongoing user) and, for ongoing users only, date of next pack start. Participants chose the time they wanted messages to be sent and could change the time by logging onto the study's web portal. Participants could also change their contact information or discontinue receiving messages at any time. We informed participants that all medical questions should be directed to their health care providers.

We collected participant contact information, including two alternate contacts, at enrollment. We planned for participants to undergo a single telephone interview 6 months after enrollment to assess our primary outcome, OCP continuation. Contact attempts began at 5 months and, for difficult-to-reach participants, continued until 8 months. To achieve desired follow-up rates, we added compensation for completing the 6-month interview. We asked participants if they had taken a pill within the last 7 days, which we defined as continuation. Because some interviews took place 6–8 months after enrollment, and thus up to 2 months after the intervention had ended, we also asked the participant if she had been taking the pill on the date 6 months after her enrollment. We further assessed continuation by asking about missed pills, interruptions in OCP use of greater than 7 days, and use of the OCP at last sexual intercourse. We created a variable for cell phone service interruptions during the study; participants were assigned to this variable if they reported either a new cell phone number or service provider or service suspensions in the past month. To have 80% power ( $P=.05$ , two-sided) to detect a 10% change in OCP continuation between the two groups, with an assumed 6-month continuation rate of 40% in the routine care group and a 15% loss to follow-up, we required 480 women in each group (960 total).

Using an intent-to-treat approach, we tested the hypothesis that educational daily text messages would affect oral contraceptive continuation. We used Student's  $t$  test to compare continuous variables and Pearson's  $\chi^2$  test to compare categorical variables when describing the population and assessing characteristics associated with 6-month OCP continuation. We conducted multivariable analyses with logistic regression using variables associated with continuation in univariable analyses. We evaluated for fit using likelihood ratio tests. We also evaluated outcomes in subgroups using both univariable and multivariable analyses. We used SPSS 18 software to compute all statistics.

## RESULTS

We screened women from March 2008 through July 2009; 78% were ineligible, 4,163 (80%) of these electing a non-OCP method of birth control ([Fig. 1](#)). Only 212 (4%) clients screened had no cell phone. We randomized 968 participants. Participants randomized to the text message intervention began receiving messages a median of 4 days (range 0–84 days) after enrollment. Four participants randomized to the intervention never received texts because they provided blocked or nonworking phone numbers. Forty-two participants randomized to the intervention discontinued text messages during the study after receiving a median of 71 messages (range 1–170).



Fig. 1

Participants were aged 13 to 25 years with 35% younger than age 20 years. Eighty-four percent were born in the United States and 60% were students. Only 11% had delivered a child, but 88% percent planned future childbearing ([Table 1](#)). Most were in a sexual relationship and satisfied with that relationship, but few were cohabiting.

Table 1

Twenty percent of participants were OCP users at enrollment (ie, took a pill within the last 7 days) and 75% ever used OCPs. Of ever OCP users, 22% discontinued past OCP use because they could not remember to take their medication. Almost all participants (98%) planned to use the pill for at least 6 months.

Seventeen percent of participants took concomitant medications, most commonly for asthma (50%), a psychiatric condition (13%), or seasonal allergies (11%). Forty percent reported forgetting and 33% worrying about forgetting to take these medications on time.

Participants reported having the same cell phone number for almost 3 years, but 13% reported at least one service suspension in the past 6 months lasting an average of 12 days. Only 2% of participants shared a phone. Participants reported sending and receiving 171 text messages per day and only 9% ever worried about the cost of text messaging. Randomization yielded similar groups, except that intervention group participants were more likely to have ever used OCPs (79% compared with 71%), more likely to have forgotten to take pills before (61% compared with 54%), and more likely to have missed two or more pills per month (34% compared with 28%) than control group participants.

We obtained follow-up information from 683 (71%) participants at a 6-month telephone interview. We made up to 31 attempts (median, four) to reach participants. The median time to follow-up was 204 days from enrollment. Of those reached, 28% reported cell phone service interruptions or changes during the study period; this did not differ by group assignment. Thirty-eight percent of loss to follow-up was the result of service interruptions of all contact numbers provided at enrollment. Compared with participants reached, those lost to follow-up were younger, had completed fewer years of school, were less likely to identify as Hispanic, more likely to have been pregnant or given birth, and less likely to be current users of OCPs (all  $P < .03$ ).

Most participants were satisfied with the number (91%), content (91%), and length (90%) of the messages received. Eighty-five percent reported the messages helped them remember to take the OCP with 81% taking a pill within 1 hour of receiving a text. Nearly half (49%) wanted to continue receiving the messages.

The intervention group ( $n=346$ ) consistently had higher OCP continuation rates than the routine care group ( $n=337$ ) regardless of how we defined continuation ([Table 2](#)). For every 10 women we enrolled in this 6-month daily text message intervention, we noted improved OCP continuation in one of them. Exploratory analyses revealed that OCP continuation was greatest in the intervention group compared with the control group (75% compared with 54%,  $P=.003$ ) if the follow-up interview occurred while the intervention was ongoing (ie, within 187 days of enrollment). Intervention group participants reached after 187 days remained somewhat more likely to continue OCPs (60% compared with 54%,  $P=.16$ ). Oral contraceptive pill continuation remained stable (52–55%) in the control group whether follow-up occurred at 5, 6, 7, or 8 or more months after enrollment. However, the improved continuation in the intervention group gradually decreased over time from 82% to 54% such that by 8 months, the effect of the intervention was extinguished.

Follow-up time point	Intervention group (%)	Control group (%)	P-value
0-187 days	75	54	.003
187-364 days	60	54	.16
364-541 days	55	52	.16
541-728 days	54	52	.16
728-915 days	54	52	.16
915-1102 days	54	52	.16
1102-1289 days	54	52	.16
1289-1476 days	54	52	.16
1476-1663 days	54	52	.16
1663-1850 days	54	52	.16
1850-2037 days	54	52	.16
2037-2224 days	54	52	.16
2224-2411 days	54	52	.16
2411-2598 days	54	52	.16
2598-2785 days	54	52	.16
2785-2972 days	54	52	.16
2972-3159 days	54	52	.16
3159-3346 days	54	52	.16
3346-3533 days	54	52	.16
3533-3720 days	54	52	.16
3720-3907 days	54	52	.16
3907-4094 days	54	52	.16
4094-4281 days	54	52	.16
4281-4468 days	54	52	.16
4468-4655 days	54	52	.16
4655-4842 days	54	52	.16
4842-5029 days	54	52	.16
5029-5216 days	54	52	.16
5216-5403 days	54	52	.16
5403-5590 days	54	52	.16
5590-5777 days	54	52	.16
5777-5964 days	54	52	.16
5964-6151 days	54	52	.16
6151-6338 days	54	52	.16
6338-6525 days	54	52	.16
6525-6712 days	54	52	.16
6712-6899 days	54	52	.16
6899-7086 days	54	52	.16
7086-7273 days	54	52	.16
7273-7460 days	54	52	.16
7460-7647 days	54	52	.16
7647-7834 days	54	52	.16
7834-8021 days	54	52	.16
8021-8208 days	54	52	.16
8208-8395 days	54	52	.16
8395-8582 days	54	52	.16
8582-8769 days	54	52	.16
8769-8956 days	54	52	.16
8956-9143 days	54	52	.16
9143-9330 days	54	52	.16
9330-9517 days	54	52	.16
9517-9704 days	54	52	.16
9704-9891 days	54	52	.16
9891-10078 days	54	52	.16

Table 2

The first interruption in pill use was 16 days later in the intervention group than the control group ( $P=.04$ ). Control group participants reported missing more pills per month (1.1 compared with 0.7,  $P=.01$ ). Ninety-four percent of participants were sexually active during the study period. At last sexual intercourse, 69% of the intervention group were OCP users compared with 60% of the control group ( $P=.03$ ). We repeated all analyses excluding the 69 women who never started the pill or were not sexually active; results for continuation were nearly identical (data not shown).

We also assessed baseline variables (shown in [Table 1](#)) to identify other factors associated

with OCP continuation in this study. [Table 3](#) lists those most associated with continuation (ie,  $P<.05$ ). Those who continued OCP use were less likely to have had cell phone service interruptions or changes during the study period (22% compared with 36%,  $P<.001$ ). Exploratory stratified analyses revealed that women with cell phone service interruptions ( $n=189$ ) were much more likely to benefit from the intervention compared with those with no interruptions (25% improvement in continuation compared with 6%,  $P<.001$ ).

Variable	Odds Ratio	P-value
Age	1.02	.001
Race	1.05	.001
Ethnicity	1.03	.001
Cell Phone Service Interruptions	0.45	<.001

Table 3

We evaluated variables that were associated with continuation in multivariable analyses. We evaluated model fit using the likelihood ratio test and present the model with best fit in [Table 4](#). After adjusting, text intervention group participants remained more likely than control participants to continue the OCPs ( $P=.03$ ). Analyses stratified by age and history of OCP use revealed no difference in the effect of the intervention. Analyses stratified by race and ethnicity showed a slightly stronger effect in African Americans and whites than in Hispanics and Asians; further analyses showed no evidence of a statistically significant interaction (data not shown).

Variable	Odds Ratio	P-value
Age	1.02	.001
Race	1.05	.001
Ethnicity	1.03	.001
Text Intervention	1.10	.03

Table 4

## DISCUSSION

This is a randomized study that evaluates the effect of text message reminders on OCP continuation. The 6-month continuation rate in the control group was similar to other recent reports.<sup>3-5</sup> Six-month OCP continuation improved by 10% in the intervention group. This is a clinically meaningful difference; we are aware of only one intervention (enhanced pill pack supply) that has achieved such success.<sup>4</sup> Furthermore, the text message reminders were also associated with OCP use at last sexual intercourse and with fewer missed pills during the trial. After the intervention ended, however, continuation rates gradually decreased to equal those in the control group.

This study included mostly young, minority women, a population with low rates of contraceptive continuation<sup>3,5</sup> and high rates of unintended pregnancy.<sup>17</sup> From a public

health perspective, we need effective strategies to help these most at-risk women reach national family planning goals.<sup>18</sup> Unlike programs that seek to change the behavior of an individual woman to increase OCP continuation (eg, through enhanced counseling), the text messaging intervention used in this study instead adapts the health system to improve outcomes. Such a strategy enables health care providers to enhance the contraceptive success of their patients simply by augmenting their clinical practice.

Limitations of this study include reliance on participant self-report of ongoing OCP use, inadvertent variable time to follow-up, and lack of tailored text message content. We found that women missed an average of one pill per month, whereas Hou, using a pill tracking device, found women missed five pills per month.<sup>13</sup> Our participants likely underreported missed pills, as has been noted elsewhere.<sup>19</sup> This may indicate socially desirable reporting, and, of more concern, that women markedly underestimate how often they miss pills and may be unaware of their consequent increased risk of pregnancy. Consistent OCP use is key to avoiding ovulation<sup>20</sup> and thus unintended pregnancy. Our exploratory analyses ([Table 2](#)) revealed that participants with timely follow-up had higher continuation rates than those who had relatively late follow-up. This may be because the effect of the intervention was limited to women currently receiving the messages. It may also be that participants who were easier to reach may represent better OCP takers because they differ in unmeasured ways from the harder-to-reach subset. This study cannot distinguish which of these possible reasons might explain the difference. Lastly, in this trial, messages were not personalized; all participants received the same messages with identical content. Tailoring message content to the individual's education level, baseline OCP knowledge, or stage of change<sup>21</sup> may increase the success of text messaging interventions.

Few interventions to improve medication adherence have shown improvements in adherence and treatment outcomes.<sup>22</sup> Many interventions have been complex and labor-intensive, limiting their usefulness in clinical settings.<sup>22</sup> In contrast, our study of a mobile health intervention used a ubiquitous device, the cell phone, and a technology, text messaging, that is increasingly used by reproductive-aged populations and can be automated to reach large populations. We previously found that most urban patients, even those without commercial insurance, have and use cell phones.<sup>23</sup> An online survey of 2,000 people similarly found that 79% of those with Medicaid regularly use their cell phones for text messaging.<sup>24</sup>

Based on these results, sending 180 daily text messages to 10 women would improve 6-month OCP continuation for one woman. To set up a similar system, health care providers would pay an upfront fee to establish the service; subsequent per-message costs are low. Furthermore, free-to-end-user systems would eliminate cost concerns for women without unlimited text plans. Clinical facilities can incorporate this technology within electronic medical record systems. Alternately, women may self-select to enroll in online or smart phone reminder systems.

Cell phone service interruptions such as suspensions, lost phones, or changes in cell phone providers were common in this study (28%). Paradoxically, exploratory analyses

demonstrated a stronger effect of the intervention in women who reported interruptions. Interruptions may reflect a busier or more disorganized life, which may, in turn, indicate a group at higher risk for OCP discontinuation. Thus, populations likely to have service interruptions may be those who benefit most from the service. Interruptions in cell phone service must be anticipated and managed but need not serve as a deterrent to implementing a similar intervention. More investigation into cell phone variables and their effect on text messaging interventions is needed.

This intervention was successful in increasing OCP continuation at 6 months and continuation was enhanced most while the intervention was ongoing. The Transtheoretical Model contends that a behavior is maintained after 6 months of consistent implementation<sup>21</sup>; however, we found that the improved continuation in the intervention group did not persist after the texts were discontinued at 6 months. Therefore, the ideal time period needed to build lasting pill-taking habits is unclear. Future studies should also evaluate whether message content is pertinent or whether a message notification alone, regardless of content, is enough to improve continuation. This intervention is ready for use in clinical practice but would benefit from large-scale evaluation.

## REFERENCES

1. Adherence to long-term therapies: evidence for action. Geneva (Switzerland): World Health Organization, 2003. Available at: <http://whqlibdoc.who.int/publications/2003/9241545992.pdf>. Retrieved August 15, 2006.
2. Mosher WD, Jones J. Use of contraception in the United States: 1982–2008. Vital and health statistics. Series 23, No. 29. Washington (DC): National Center for Health Statistics; 2010. DHHS publication no. [PHS] 2010-1981.
3. Nelson AL, Westhoff C, Schnare SM. Real-world patterns of prescription refills for branded hormonal contraceptives: a reflection of contraceptive discontinuation. *Obstet Gynecol* 2008;112:782–7.
4. White KO, Westhoff C. The effect of pack supply on oral contraceptive pill continuation: a randomized controlled trial. *Obstet Gynecol* 2011;118:615–22.
5. Westhoff C, Heartwell S, Edwards S, Ziemann M, Cushman L, Robilotto C, et al.. Initiation of oral contraceptives using a quick start compared with a conventional start: a randomized controlled trial. *Obstet Gynecol* 2007;109:1270–6.
6. Rosenberg MJ, Waugh MS, Long S. Unintended pregnancies and use, misuse and discontinuation of oral contraceptives. *J Reprod Med* 1995;40:355–60.
7. Rosenberg MJ, Waugh MS, Meehan TE. Use and misuse of oral contraceptives: risk indicators for poor pill taking and discontinuation. *Contraception* 1995;51:283–8.

8. Free C, Knight R, Robertson S, Whittaker R, Edwards P, Zhou W, et al.. Smoking cessation support delivered via mobile phone text messaging (txt2stop): a single-blind, randomised trial. *Lancet* 2011;378:49–55.
9. Rodgers A, Corbett T, Bramley D, Riddell T, Wills M, Lin R-B, et al.. Do u smoke after txt? results of a randomized trial of smoking cessation using mobile phone text messaging. *Tob Control* 2005;14:255–61.
10. Menon-Johansson AS, McNaught F, Mandalia S, Sullivan AK. Texting decreases the time to treatment for genital Chlamydia trachomatis infection. *Sex Transm Infect* 2006;82:49–51.
11. Miloh T, Annunziato R, Arnon R, Warshaw J, Parkar S, Suchy FJ, et al.. Improved adherence and outcomes for pediatric liver transplant recipients using text messaging. *Pediatrics* 2009;124:e844–50.
12. Vilella A, Bayas J-M, Diaz M-T, Guinovart C, Diez C, Simó D, et al.. The role of mobile phones in improving vaccination rates in travelers. *Prev Med* 2004;38:503–9.
13. Hou MY, Hurwitz S, Kavanagh E, Fortin J, Goldberg AB. Using daily text-message reminders to improve adherence with oral contraceptives: a randomized controlled trial. *Obstet Gynecol* 2010;116:633–40.
14. Lenhart A, Ling R, Campbell S, Purcell K. Teens and mobile phones: text messaging explodes as teens embrace it as the centerpiece of their communication strategies with friends. Washington (DC): Pew Research Center; 2010. Available at: <http://pewinternet.org/Reports/2010/Teens-and-Mobile-Phones.aspx>. Retrieved April 21, 2010.
15. US teen mobile report: calling yesterday, texting today, using apps tomorrow. The Nielsen Company; 2010. Available at: [http://blog.nielsen.com/nielsenwire/online\\_mobile/u-s-teen-mobile-report-calling-yesterday-texting-today-using-apps-tomorrow/](http://blog.nielsen.com/nielsenwire/online_mobile/u-s-teen-mobile-report-calling-yesterday-texting-today-using-apps-tomorrow/). Retrieved October 15, 2010.
16. Hatcher RA, Trussell T, Stewart F, Nelson AL, Cates W, Guest F, et al.. Contraceptive technology. 18th revised ed. New York (NY): Ardent Media; 2004.
17. Finer LB, Henshaw SK. Disparities in rates of unintended pregnancy in the United States, 1994 and 2001. *Perspect Sex Reprod Health* 2006;38:90–6.
18. Healthy People 2020 summary of objectives: family planning. Washington (DC): US Department of Health and Human Services; 2001. Available at: <http://healthypeople.gov/2020/topicsobjectives2020/pdfs/FamilyPlanning.pdf>. Retrieved July 2, 2011.
19. Potter L, Oakley D, de Leon-Wong E, Cañamar R. Measuring compliance among oral

contraceptive users. *Fam Plann Perspect* 1996;28:154–8.

20. Westhoff CL, Torgal AH, Mayeda ER, Stanczyk FZ, Lerner J, Benn EKT, et al.. Ovarian suppression in normal-weight and obese women during oral contraceptive use. *Obstet Gynecol* 2010;116:275–83.

21. Johnson SS, Grimley DM, Prochaska JO. Prediction of adherence using the transtheoretical model: implications for pharmacy care practice. *J Soc Adm Pharm* 1998;15:135–48.

22. Haynes RB, McKibbin KA, Kanani R. Systematic review of randomized trials of interventions to assist patients to follow prescriptions for medications. *Lancet* 1996;348:383–6. Erratum in *Lancet* 1997;349:1180.

23. Castaño P, Martínez RA. Harnessing technology for adolescent health promotion. *Adolesc Med State Art Rev* 2007;18:400–6, xiii.

24. Healthcare unwired: new business models delivering care anywhere. PricewaterhouseCoopers' Health Research Institute, 2010. Available at: <http://pwchealth.com/cgi-local/hregister.cgi?link=reg/healthcare-unwired.pdf>. Retrieved February 9, 2011.

### Supplemental Digital Content

- [AOG 119 1 2011 12 06 CASTANO 202309 SDC1.pdf; \[PDF\] \(15 KB\)](#)

© 2012 The American College of Obstetricians and Gynecologists